Patient Enrollment



RELIZORB should only be used in conjunction with an enteral feeding system that has a low-flow/no-flow alarm (pump rate should be set between 24-120 mL/hour). RELiZORB should not be used with formulas that contain insoluble fiber. For more information regarding RELiZORB use, visit www.relizorb.com, or call 1-844-632-9271.

Please complete this form and email to info@relizorbsupport.com or fax to 1-844-233-3146. Please note—ALL INFORMATION IS REQUIRED to expedite processing of referral.

1. Patient Information							
Name (First):	_(Last):						
Street Address:							
SSN #:							
Patient/Patient Representative C	Contact Information:						
	Patient Representative Name:						
	Patient Representative Relationship::						
2. Current Insurance Info	rmation						
Primary Insurance Plan:	☐ Private/Commercial ☐ Medicaid	☐ Medicare ☐ Medicare Advantage	\square Patient has no insurance				
Primary Insurance Name:							
Insurance Phone #:	Member ID #:						
Policy Holder:	Policy Holder Date of Birth: Relationship to Patient:						
Secondary Insurance Plan:	☐ Private/Commercial ☐ Medicaid	☐ Medicare ☐ Medicare Advantage	☐ Patient ha	s no insurance			
Secondary Insurance Name		_					
	Policy Holder Date of Birth: Relationship to Patient:						
NOTE: Please attach a copy of th	e insurance card (front and ba	CK)					
3. Prescriber Information							
Proscriber Name (First):		(Last):					
NPI:		(Last):					
Tax ID #:	PTAN:						
Prescriber Specialty:							
Center/Hospital Name:				_			
			State	: ZIP:			
Prescriber Direct Contact #:							
Best day(s)/times if peer-to-peer							
Primary Office Contact:				Ext:			
Email:	Fax:						

4. Prescription for REL	iZORB® (iMMOBILIZ	ZED LIPASE) (ARTRIDGE				
In order for us to send RELiZ	ORB to your patient, the	prescription info	rmation must be	e complete and a	accurate.		
Patient Name (First, Last):					Date of Birth:		
Primary Diagnosis Code:							
Other Diagnosis Code(s):							
Height:					□ lb □kg		
					cement Date:		
					e (mL/hour):		
Product Name: RELiZORB® (i							
RELIZORB PRESCRIPTION (constructions: Use 1 cartridge in used/day)		g tube set, change	cartridge with e	every 500 mL of	enteral formula (max of 2 cartridges		
□1 cartridge/day (500 mL) Dispense 30 each/cartrid		□ 2 cartridges/da Dispense 60 ea			☐ No. of refills:		
Additional Orders/Comment	s:						
I certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.							
Doctor/Prescriber Signature	>				Date:		
5. Continuity of Care/H	lospital Discharge						
RELIZORB is committed to your patient's continuity of care on the journey to home.							
☐ Please check this box for hospital discharge patients							
6. Clinical Indications	Supporting Medical	Necessity: (P	lease check	all that apply			
☐ Patient has failed to achie		·	- '	t therapy in conju	unction with enteral feeding.		
☐ Diarrhea	☐ Fatty stools	Abdomin	al pain	□ Bloating			
□ Nausea	\square Constipation	☐ Flatulence	е	□Vomiting			
□ Patient demonstrates failure to achieve or maintain target BMI. □ Patient requires overnight enteral feeding to meet caloric and nutritional demands with need for sustained lipase delivery throughout feed. □ Patient exhibits deficiency in fatty acid levels. □ Patient's symptoms of fat malabsorption impair or inhibit patient's activites of daily living and quality of life.							
Doctor/Prescriber Signature	>				Date:		
7. Please Include the F	ollowing Clinical Do	cumentation					
☐ Copy of front and back of ☐ MD office visit notes inclu		P, referrals	☐ RD office not☐ Medication lis		☐ Weight history ☐ Letter of medical necessity, if needed		

